

Sinle-valued and spinor representations of orientation-preserved four-dimensional cubic group

Jian Dai*, Xing-Chang Song†
Department of Physics, Peking University

February 23, 1999
Revised on September 6, 2000

Abstract

We calculate the single-valued and spinor representations of SO_4 , the orientation-preserved subgroup of O_4 , on the base of our previous work on four-dimensional cubic group O_4 ,

*daij1492@yahoo.com

†songxc@ibm320h.phy.pku.edu.cn

I Introduction

This paper should be considered as a continuation of our previous work [1] where the single-valued representations and spinor representations of four-dimensional cubic group O_4 are obtained. We will consider here the orientation-preserved subgroup of O_4 , namely SO_4 , and derive the representation theory of its double group $\overline{SO_4}$. The representation theories of SO_4 and O_4 are well-known results for years [2][3]; we re-derive them adopting a more systematic method and give their spinor parts which are totally new. This paper is arranged as following. First we construct SO_4 , extend it to double group $\overline{SO_4}$ and calculate their conjugate classes. Then we reduce all single-valued representations as well as two-valued representations of SO_4 from those of O_4 based on our previous work. From the point of physical application, we expect that fermions in lattice field theory should transform according to spinor representations of symmetry group of the lattice where they reside.

II Structure of $\overline{SO_4}$

For completeness, we recall some definitions and results in [1]. An n -cube (or *hyper-cube in E^n*) C_n is defined to be a subset of E^n , $C_n = \{p|x^i(p) = \pm 1\}$, where $x^i : E^n \rightarrow \mathcal{R}, i = 1, 2, \dots, n$ are coordinate functions of E^n , together with the distance inherited from E^n . n -Cubic group (*hyper-cubic group of degree n*) O_n consists of all isometries of E^n which stabilize C_n and has a structure of wreath product

$$O_n \cong Z_2^n \rtimes S_n \quad (1)$$

in which the first factor Z_2^n is generated by the inversions of axes and the second factor S_n is the permutations of axes. The wreath product structure (1) enable us to regard O_n as a permutation group with signatures whose elements can be decomposed into independent cycles with signature. A k -cycle with signature in $Z_2^n \rtimes S_n$ is represented as

$$\left(\begin{array}{cccc} a_1 & a_2 & \dots & a_k \\ (-)^{s_{a_1}} a_2 & (-)^{s_{a_2}} a_3 & \dots & (-)^{s_{a_k}} a_1 \end{array} \right), s_{a_i} \in \mathcal{Z}/2\mathcal{Z}, i = 1, 2, \dots, k$$

whose signature is defined to be $(-)^{s_{a_1} + s_{a_2} + \dots + s_{a_k}}$. A conjugate classes of O_n can be uniquely determined by the cycle structure descended from S_n and signatures of its cycles [4].

Let $n = 4$ below. O_4 is generated by a set of generators $\{e_i, i = 1, 2, 3, 4; \gamma, t\}$ together with a system of constraint equations

$$e_i^2 = e, e_i e_j = e_j e_i, i, j = 1..4, i \neq j \quad (2)$$

$$\gamma^2 = e, t^3 = e, (t\gamma)^4 = e \quad (3)$$

$$te_1 = e_1, te_2 = e_4 t, te_3 = e_2 t, te_4 = e_3 t \quad (4)$$

$$\gamma e_1 = e_3 \gamma, \gamma e_2 = e_2 \gamma, \gamma e_4 = e_4 \gamma \quad (5)$$

in which e_i s generate Z_2^4 and γ, t generate S_4 . All of 20 conjugate classes of O_4 can be depicted by a class of generalized Young diagrams where a slash inside a Young box shows that there is a minus sign in the position of corresponding cycle (see table I).

We know from the theory of Clifford algebra [5] that there is a short exact sequence

$$0 \rightarrow Z_2 \rightarrow Pin(E^n) \xrightarrow{\pi} O(n) \rightarrow 1$$

Let ϵ be an injective homomorphism from a group G to $O(n)$, then the double group or the spin-extension of G with respect to ϵ is defined to be $D_n(G, \epsilon) := \pi^{-1}(G)$. With n and ϵ fixed, we use $\tilde{G}$ to denote $D_n(G, \epsilon)$. Let C be a conjugate class in G , then either will $\pi^{-1}(C)$ be one conjugate class in $\tilde{G}$ still or it will split into two conjugate classes in $\tilde{G}$ containing $s(g), -s(g), \forall g \in C$ respectively, where $s : G \rightarrow \tilde{G}, s.t. \pi s = Id_G$.

$\overline{O_4}$ is generated by lifting Eq.(2)(3)(4)(5) to

$$e_i^2 = -1, e_i e_j = -e_j e_i, i, j = 1..4, i \neq j \quad (6)$$

$$\gamma^2 = -1, t^3 = -1, (t\gamma)^4 = -1 \quad (7)$$

$$te_1 = e_1 t, te_2 = -e_4 t, te_3 = -e_2 t, te_4 = e_3 t \quad (8)$$

$$\gamma e_1 = -e_3 \gamma, \gamma e_2 = -e_2 \gamma, \gamma e_4 = -e_4 \gamma \quad (9)$$

Each conjugate class of O_4 is lifted into one corresponding conjugate class in $\overline{O_4}$, except classes 1, 8, 14, 15, 20 which splits into two classes each of them. Hence there are total 25 classes of $\overline{O_4}$.

The *orientation-preserved n-cubic group* is defined to be

$$SO_n := (O_n \cap SO(n)) \triangleleft O_n$$

The generators of Z_2^n are written as $e_i, i = 1, 2, \dots, n$. Define $Z_2^n|_e$ as a subgroup of Z_2^n generated by $e_i e_j, i \neq j$ and $Z_2^n|_o := Z_2^n \setminus Z_2^n|_e$. Then

$$SO_n = (Z_2^n|_e \cdot A_n) \bigsqcup (Z_2^n|_o \cdot (S_n \setminus A_n)) \quad (10)$$

in which $\cdot$ is product of two subsets in a group, A_n stands for the alternative subgroup in S_n . Thus, $|SO_n| = (2^n \cdot (n!))/2$.

Specify $n = 4$ and we know immediately that $|SO_4| = 192$. We introduce

$$\eta = \gamma t \gamma t^2 \gamma; \quad (11)$$

$$\alpha = (t^2 \gamma)^2, \beta = t \gamma t^2 \gamma t; \quad (12)$$

$$x = e_1 \eta, y = e_4 \eta, q = e_2 \eta \quad (13)$$

then the structure of SO_4 can be summarized as

$$x^2 = y^2 = q^4 = e, yx = xy, qx = xq^3, qy = yq^3 \quad (14)$$

$$\alpha^2 = \beta^2 = t^3 = e, \beta\alpha = \alpha\beta, t\alpha = \alpha\beta t, t\beta = \alpha t \quad (15)$$

$$\alpha x = q\beta, \alpha y = q^3\beta, \alpha q = x\beta \quad (16)$$

$$\beta x = q^3\alpha, \beta y = q\alpha, \beta q = y\alpha \quad (17)$$

$$tx = xt^2, ty = q^3 t^2, tq = yt^2 \quad (18)$$

together with

$$x\alpha = \beta q^3, xt = t^2 x; y\beta = \alpha q^3, yt = t^2 q^3; qt = t^2 y \quad (19)$$

Accordingly, each group element can be expressed as a "normal ordering" product of $x, y \rightarrow q \rightarrow \alpha, \beta \rightarrow t$ and their powers from left to right. Throwing away all classes which belong to O_4 but not to SO_4 , there are 11 left which are 1, 3, 5, 7, 8, 11, 12, 14, 15, 18, 20 in Table I. The 14th and the 20th will part into two classes with equal numbers of elements each under adjoint action of SO_4 which are denoted as 14, 14', 20, 20'. Therefore, there are 13 conjugate classes in SO_4 .

Due to the fact that $\overline{SO_4} \triangleleft \overline{O_4}$ and theory of group extension [6], the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & Z_2 & \longrightarrow & \overline{O_4} & \xrightarrow{\pi} & O_4 \longrightarrow 1 \\ & & \parallel & & \uparrow & & \uparrow \\ 0 & \longrightarrow & Z_2 & \longrightarrow & \overline{SO_4} & \longrightarrow & SO_4 \longrightarrow 1 \end{array}$$

is commutative. We can lift generating relations (14)(15)...(19) to

$$\begin{aligned}
x^2 &= y^2 = q^4 = -1, yx = -xy, qx = -xq^3, qy = -yq^3 \\
\alpha^2 &= \beta^2 = t^3 = -1, \beta\alpha = -\alpha\beta, t\alpha = -\alpha\beta t, t\beta = \alpha t \\
\alpha x &= q\beta, \alpha y = q^3\beta, \alpha q = x\beta \\
\beta x &= q^3\alpha, \beta y = q\alpha, \beta q = y\alpha \\
tx &= -xt^2, ty = q^3t^2, tq = yt^2 \\
x\alpha &= \beta q^3, xt = -t^2x; y\beta = \alpha q^3, yt = t^2q^3; qt = t^2y
\end{aligned}$$

by the definition (11)(12)(13). As to the conjugate classification, 1, 5, 8, 11, 14, 15, 18, 20, 14', 20' violate relation $-g \sim g$, so that $\overline{SO_4}$ is partitioned into 23 classes, suggesting that there are altogether 23 inequivalent irreducible representations in which 13 representations are single-valued to SO_4 . We summarize the conjugate classes of $\overline{SO_4}$ in Table II.

III Representation theory of $\overline{SO_4}$

All inequivalent irreducible representations of $\overline{SO_4}$ can be reduced out from those of $\overline{O_4}$ [1]. Table III gives the characters of all inequivalent irreducible representations of $\overline{O_4}$, respect to the classes of $\overline{SO_4}$. A brief observation gives some important information. Firstly, $1_1 \cong 1_4, 1_2 \cong 1_3, 2_1 \cong 2_2, 3_1 \cong 3_4, 3_2 \cong 3_3, 4_1 \cong 4_4, 4_2 \cong 4_3, 8_1 \cong 8_2, 6_1 \cong 6_3$. Secondly, omitting equivalence, $1_1, 1_2, 2_1, 3_1, 3_1, 4_1, 4_2, 8_1, 6_1$, remain irreducible within $\overline{SO_4}$, while other 7 become reducible. Thirdly, as for each of these reducible ones, the inner product of itself's character equals to 2, implying that it can be reduced to two inequivalent irreducible representations, thus there are 13 single-valued and 10 spinor representations as we expected. Finally, it is one possible solution to Burside theorem that each of these seven reducible representations splits into 2 inequivalent irreducible representations with equal dimensions. We conjecture that it is the solution to our representation theory of $\overline{SO_4}$ and try to verify it below.

Summarily speaking, there are nine single-valued inequivalent irreducible representations inherited from $\overline{O_4}$

$$1_1, 1_2, 2 \equiv 2_1, 3_1, 3_2, 4_1, 4_2, 8 \equiv 8_1, 6 \equiv 6_1$$

and we conjecture the splitting relations

$$\begin{aligned}
6_2, 6_4 &\rightarrow 3_\alpha, 3_\beta, 3_\gamma, 3_\delta \\
4_1, 4_2 &\rightarrow 2_\alpha, 2_\beta, 2_\gamma, 2_\delta \\
8 &\rightarrow 4_\alpha, 4_\beta \\
12_1, 12_2 &\rightarrow 6_\alpha, 6_\beta, 6_\gamma, 6_\delta
\end{aligned}$$

III.1 Hidden single-valued representations

The representation matrices of x, y, q in 6_2 [1] are written as

$$x \mapsto \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & -1 & 0 \end{pmatrix}, y \mapsto \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 \end{pmatrix}, q \mapsto \begin{pmatrix} -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & -1 & 0 \end{pmatrix}$$

The textures of these matrices inspire us to such a hypotheses that in $3_{\alpha,\beta,\gamma,\delta}$, x, y, q take on a form like

$$x, y \mapsto \begin{pmatrix} \pm 1 & \\ & \pm H \end{pmatrix}, q \mapsto \begin{pmatrix} \pm 1 & \\ & \pm Q \end{pmatrix}$$

where $H \equiv \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, $Q \equiv \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. After taking account of the conjugate equivalence, only four possibilities survive from the totally 64, i.e.

$$I : x \rightarrow \begin{pmatrix} 1 & \\ & H \end{pmatrix}, y \rightarrow \begin{pmatrix} 1 & \\ & -H \end{pmatrix}, q \rightarrow \begin{pmatrix} -1 & \\ & Q \end{pmatrix} \quad (20)$$

$$II : x \rightarrow \begin{pmatrix} -1 & \\ & H \end{pmatrix}, y \rightarrow \begin{pmatrix} -1 & \\ & -H \end{pmatrix}, q \rightarrow \begin{pmatrix} 1 & \\ & Q \end{pmatrix} \quad (21)$$

$$III : x \rightarrow \begin{pmatrix} 1 & \\ & -H \end{pmatrix}, y \rightarrow \begin{pmatrix} 1 & \\ & H \end{pmatrix}, q \rightarrow \begin{pmatrix} -1 & \\ & Q \end{pmatrix} \quad (22)$$

$$IV : x \rightarrow \begin{pmatrix} -1 & \\ & -H \end{pmatrix}, y \rightarrow \begin{pmatrix} -1 & \\ & H \end{pmatrix}, q \rightarrow \begin{pmatrix} 1 & \\ & Q \end{pmatrix} \quad (23)$$

which also satisfy Eq.(14). Then we regard α, β, t as unknowns, (15)–(19) as constraint, and solve these matrix equations. Modulo similarity, each of Eqs.(20)(21) (22)(23) gives two solutions, labeled as $I, I', II, II', III, III', IV, IV'$; however, there is no difficulty to find out that $I \cong III$, $I' \cong III'$, $II \cong IV$, $II' \cong IV'$. Thus I, I', II, II' are what we need.

$$3_\alpha \equiv I : \alpha \rightarrow \text{diag}(-1, 1, -1), \beta \rightarrow \text{diag}(1, -1, -1), t \rightarrow \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$$

$$3_\beta \equiv I' : \alpha \rightarrow \text{diag}(1, -1, -1), \beta \rightarrow \text{diag}(-1, -1, 1), t \rightarrow \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$$

$$3_\gamma \equiv II : \alpha \rightarrow \text{diag}(-1, 1, -1), \beta \rightarrow \text{diag}(1, -1, -1), t \rightarrow \begin{pmatrix} 0 & 0 & 1 \\ -1 & 0 & 0 \\ 0 & -1 & 0 \end{pmatrix}$$

$$3_\delta \equiv II' : \alpha \rightarrow \text{diag}(1, -1, -1), \beta \rightarrow \text{diag}(-1, -1, 1), t \rightarrow \begin{pmatrix} 0 & 0 & 1 \\ -1 & 0 & 0 \\ 0 & -1 & 0 \end{pmatrix}$$

III.2 Spinor representations

It is more straightforward to reduce out the spinor representations. A deep result of Clifford theory on decomposition of induced representations [7][1] shows that the spinor representation matrices of O_4 are of the form of a tensor product

$$S_i(g) = S(g) \otimes s_i(g), \forall g \in \overline{O_4}, i = \underline{4}_1, \underline{4}_2, 8, 12_1, 12_2$$

where S is given by the algebraic isomorphism from $Cl(E^4)$ to $M_2(\mathbf{H})$ and s_i has the same texture (zero matrix element positions) of irreducible representation i of S_4 . Additionally, for g in $\overline{SO_4}$, $S(g)$ takes on a 2-by-2 block diagonal form

$$S(g) = \begin{pmatrix} S_{up}(g) & 0 \\ 0 & S_{down}(g) \end{pmatrix} \quad (24)$$

So it is just what we want

$$S_{up}(x) = \frac{1}{\sqrt{2}} \cdot \begin{pmatrix} e^{i(\frac{3}{4})\pi} & e^{i(\frac{-3}{4})\pi} \\ e^{i(\frac{1}{4})\pi} & e^{i(\frac{-1}{4})\pi} \end{pmatrix}, S_{up}(y) = \frac{1}{\sqrt{2}} \cdot \begin{pmatrix} e^{i(\frac{3}{4})\pi} & e^{i(\frac{1}{4})\pi} \\ e^{i(\frac{-3}{4})\pi} & e^{i(\frac{-1}{4})\pi} \end{pmatrix} \quad (25)$$

$$S_{up}(q) = \frac{1}{\sqrt{2}} \cdot \begin{pmatrix} e^{i(\frac{1}{4})\pi} & e^{i(\frac{3}{4})\pi} \\ e^{i(\frac{3}{4})\pi} & e^{i(\frac{1}{4})\pi} \end{pmatrix}, S_{up}(t) = \frac{1}{\sqrt{2}} \cdot \begin{pmatrix} 1 & -i \\ 1 & i \end{pmatrix} \quad (26)$$

$$S_{up}(\alpha) = \begin{pmatrix} 0 & -i \\ -i & 0 \end{pmatrix}, S_{up}(\beta) = \begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix} \quad (27)$$

$$S_{down}(x) = \frac{1}{\sqrt{2}} \cdot \begin{pmatrix} e^{i(\frac{3}{4})\pi} & e^{i(\frac{-3}{4})\pi} \\ e^{i(\frac{1}{4})\pi} & e^{i(\frac{-1}{4})\pi} \end{pmatrix}, S_{down}(y) \rightarrow \frac{1}{\sqrt{2}} \cdot \begin{pmatrix} e^{i(\frac{-1}{4})\pi} & e^{i(\frac{-3}{4})\pi} \\ e^{i(\frac{1}{4})\pi} & e^{i(\frac{3}{4})\pi} \end{pmatrix} \quad (28)$$

$$S_{down}(q) = \frac{1}{\sqrt{2}} \cdot \begin{pmatrix} e^{i(\frac{1}{4})\pi} & e^{i(\frac{3}{4})\pi} \\ e^{i(\frac{3}{4})\pi} & e^{i(\frac{1}{4})\pi} \end{pmatrix}, S_{down}(t) = \frac{1}{\sqrt{2}} \cdot \begin{pmatrix} i & 1 \\ -i & 1 \end{pmatrix} \quad (29)$$

$$S_{down}(\alpha) = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, S_{down}(\beta) = \begin{pmatrix} 0 & -i \\ -i & 0 \end{pmatrix} \quad (30)$$

Keeping the second factor unchanged, each spinor representation in $\overline{O_4}$ splits into two spinor representations in $\overline{SO_4}$, denoted as $2_\alpha, 2_\beta, 2_\gamma, 2_\delta, 4_\alpha, 4_\beta, 6_\alpha, 6_\beta, 6_\gamma, 6_\delta$.

In fact, $S(g)$ falls in the so-called "chiral"-representation of $Cl(E^4)$ in physical language. Due to $Cl(V) = Cl(V)_e \oplus Cl(V)_o$, and the choice of chiral-representation, there are

$$Cl(E^4)_e \cong \begin{pmatrix} \mathbf{H} & O \\ 0 & \mathbf{H} \end{pmatrix}, Cl(E^4)_o \cong \begin{pmatrix} 0 & \mathbf{H} \\ \mathbf{H} & 0 \end{pmatrix}$$

Notice that $\overline{SO_4} < Spin(4) \subset Cl(E^4)_e$, so our reducing process for spinor representations roots in the structure of Clifford algebra.

III.3 Results

Conclusively, our conjecture really gives all inequivalent irreducible representations whose characters are summarized in Table IV.

Acknowledgements

This work was supported by Climb-Up (Pan Deng) Project of Department of Science and Technology in China, Chinese National Science Foundation and Doctoral Programme Foundation of Institution of Higher Education in China. One of the authors JD is grateful to Dr. L-G. Jin for his advice on this paper.

References

- [1] J.Dai,X-C.Song, "Structure and representation theory of the double group of four-dimensional cubic group" (submitted to J. Math. Phy.)
- [2] M.Baake,B.Gemünden and R.Oedingen, "Structure and representations of the symmetry group of the four-dimension cube", J. Math. Phys. **23**(1982)944-953
- [3] J.E.Mandula,G.Zweig and J.Govaerts, "Representations of the rotation reflection symmetry group of the four-dimensional cubic lattice", Nucl.Phys.B**228**(1983)91-108
- [4] Adalbert Kerber, *Representations of Permutation Groups I*, Springer-Verlag (1971)
- [5] F.Reese Harvey, *Spinors and Calibrations*, academic press (1990)
- [6] Kenneth S.Brown, *Cohomology of Groups*, springer-verlag (1982)
- [7] Charles W.Curtis and Irving Reiner, *Methods of Representation Theory* vol.I, John Wiley & Sons Inc. (1981)

Table I: Conjugate Classes of $Z_2^4 \rtimes S_4$

No	SplitNo	YoungDiagram	ord	num	det	No	SplitNo	YoungDiagram	ord	num	det
1	1-1		1	1	1	2	1-2		2	4	-1
3	1-3		2	6	1	4	1-4		2	4	-1
5	1-5		2	1	1						
6	2-1		2	12	-1	7	2-2		2	24	1
8	2-3		4	12	1	9	2-4		2	12	-1
10	2-5		4	24	-1	11	2-6		4	12	1
12	3-1		2	12	1	13	3-2		4	24	-1
14	3-3		4	12	1						
15	4-1		3	32	1	16	4-2		6	32	-1
17	4-3		6	32	-1	18	4-4		6	32	1
19	5-1		4	48	-1	20	5-2		8	48	1

"SplitNo" of each class reflects the relation between the classes of $Z_2^4 \rtimes S_4$ and those of S_4 . "ord" means order of each class. "num" is the number of elements in each class. "det" is the signature of each class.

 Table II: Conjugate Classes of $\overline{SO_4}$

No.	1	1	3	5	5	7	8	8	11	11	12	14	14	14'	14'	15	15	18	18	20	20	20'	20'
num.	1	1	12	1	1	48	12	12	12	12	24	6	6	6	6	32	32	32	32	24	24	24	24
ord.	1	2	4	2	2	4	8	8	8	8	4	4	4	4	4	6	3	6	6	8	8	8	8

Table III: Character Table of $\overline{O_4}$ (with respect to the classes of $\overline{SO_4}$)

	1	1	3	5	5	7	8	8	11	11	12	14	14	14'	14'	15	15	18	18	20	20	20'	20'	(χ, χ)
$\chi_1^{(1)}$	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
$\chi_2^{(1)}$	1	1	1	1	1	-1	-1	-1	-1	-1	1	1	1	1	1	1	1	1	1	-1	-1	-1	-1	1
$\chi_1^{(2)}$	2	2	2	2	2	0	0	0	0	0	2	2	2	2	2	-1	-1	-1	-1	0	0	0	0	1
$\chi_1^{(3)}$	3	3	3	3	3	1	1	1	1	1	-1	-1	-1	-1	-1	0	0	0	0	-1	-1	-1	-1	1
$\chi_2^{(3)}$	3	3	3	3	3	-1	-1	-1	-1	-1	-1	-1	-1	-1	-1	0	0	0	0	1	1	1	1	1
$\chi_3^{(1)}$	1	1	1	1	1	-1	-1	-1	-1	-1	1	1	1	1	1	1	1	1	1	-1	-1	-1	-1	1
$\chi_4^{(1)}$	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
$\chi_2^{(2)}$	2	2	2	2	2	0	0	0	0	0	2	2	2	2	2	-1	-1	-1	-1	0	0	0	0	1
$\chi_3^{(3)}$	3	3	3	3	3	-1	-1	-1	-1	-1	-1	-1	-1	-1	-1	0	0	0	0	1	1	1	1	1
$\chi_4^{(3)}$	3	3	3	3	3	1	1	1	1	1	-1	-1	-1	-1	-1	0	0	0	0	-1	-1	-1	-1	1
$\chi_1^{(4)}$	4	4	0	-4	-4	0	2	2	-2	-2	0	0	0	0	0	1	1	-1	-1	0	0	0	0	1
$\chi_2^{(4)}$	4	4	0	-4	-4	0	-2	-2	2	2	0	0	0	0	0	1	1	-1	-1	0	0	0	0	1
$\chi_1^{(8)}$	8	8	0	-8	-8	0	0	0	0	0	0	0	0	0	0	-1	-1	1	1	0	0	0	0	1
$\chi_3^{(4)}$	4	4	0	-4	-4	0	-2	-2	2	2	0	0	0	0	0	1	1	-1	-1	0	0	0	0	1
$\chi_4^{(4)}$	4	4	0	-4	-4	0	2	2	-2	-2	0	0	0	0	0	1	1	-1	-1	0	0	0	0	1
$\chi_2^{(8)}$	8	8	0	-8	-8	0	0	0	0	0	0	0	0	0	0	-1	-1	1	1	0	0	0	0	1
$\chi_1^{(6)}$	6	6	-2	6	6	0	0	0	0	0	2	-2	-2	-2	-2	0	0	0	0	0	0	0	0	1
$\chi_2^{(6)}$	6	6	-2	6	6	2	-2	-2	-2	-2	2	2	2	2	2	0	0	0	0	0	0	0	0	2
$\chi_3^{(6)}$	6	6	-2	6	6	0	0	0	0	0	2	-2	-2	-2	-2	0	0	0	0	0	0	0	0	1
$\chi_4^{(6)}$	6	6	-2	6	6	-2	2	2	2	2	-2	2	2	2	2	0	0	0	0	0	0	0	0	2
$\chi_1^{(4)}$	4	-4	0	0	0	0	$-2\sqrt{2}$	$2\sqrt{2}$	0	0	0	2	-2	2	-2	2	-2	0	0	$\sqrt{2}$	$-\sqrt{2}$	$\sqrt{2}$	$-\sqrt{2}$	2
$\chi_2^{(4)}$	4	-4	0	0	0	0	$2\sqrt{2}$	$-2\sqrt{2}$	0	0	0	2	-2	2	-2	2	-2	0	0	$-\sqrt{2}$	$\sqrt{2}$	$-\sqrt{2}$	$\sqrt{2}$	2
$\chi^{(8)}$	8	-8	0	0	0	0	0	0	0	0	4	-4	4	-4	-2	2	0	0	0	0	0	0	0	2
$\chi_1^{(12)}$	12	-12	0	0	0	0	$-2\sqrt{2}$	$2\sqrt{2}$	0	0	0	-2	2	-2	2	0	0	0	0	$-\sqrt{2}$	$\sqrt{2}$	$-\sqrt{2}$	$\sqrt{2}$	2
$\chi_2^{(12)}$	12	-12	0	0	0	0	$2\sqrt{2}$	$-2\sqrt{2}$	0	0	0	-2	2	-2	2	0	0	0	0	$\sqrt{2}$	$-\sqrt{2}$	$\sqrt{2}$	$-\sqrt{2}$	2

Character is labeled by a superscript showing its dimension where an underline shows a spinor representation and a subscript distinguishing different representations with same dimension. (χ, χ) evaluates the inner product of a character with itself.

Table IV: Character Table of $\overline{SO}_4$

	1	$\bar{1}$	3	5	$\bar{5}$	7	8	$\bar{8}$	11	$\bar{11}$	12	14	$\bar{14}$	14'	14'	15	$\bar{15}$	18	$\bar{18}$	20	$\bar{20}$	20'	20'
num.	1	1	12	1	1	48	12	12	12	12	24	6	6	6	6	32	32	32	32	24	24	24	24
$\chi_1^{(1)}$	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
$\chi_2^{(1)}$	1	1	1	1	1	-1	-1	-1	-1	-1	1	1	1	1	1	1	1	1	1	-1	-1	-1	-1
$\chi^{(2)}$	2	2	2	2	2	0	0	0	0	0	2	2	2	2	2	-1	-1	-1	-1	0	0	0	0
$\chi_1^{(3)}$	3	3	3	3	3	1	1	1	1	1	-1	-1	-1	-1	-1	0	0	0	0	-1	-1	-1	-1
$\chi_2^{(3)}$	3	3	3	3	3	-1	-1	-1	-1	-1	-1	-1	-1	-1	-1	0	0	0	0	1	1	1	1
$\chi_1^{(4)}$	4	4	0	-4	-4	0	2	2	-2	-2	0	0	0	0	0	1	1	-1	-1	0	0	0	0
$\chi_2^{(4)}$	4	4	0	-4	-4	0	-2	-2	2	2	0	0	0	0	0	1	1	-1	-1	0	0	0	0
$\chi^{(8)}$	8	8	0	-8	-8	0	0	0	0	0	0	0	0	0	0	-1	-1	1	1	0	0	0	0
$\chi^{(6)}$	6	6	-2	6	6	0	0	0	0	0	2	-2	-2	-2	-2	0	0	0	0	0	0	0	0
$\chi_\alpha^{(3)}$	3	3	-1	3	3	1	-1	-1	-1	-1	-1	-1	-1	3	3	0	0	0	0	-1	-1	1	1
$\chi_\beta^{(3)}$	3	3	-1	3	3	1	-1	-1	-1	-1	-1	3	3	-1	-1	0	0	0	0	1	1	-1	-1
$\chi_\gamma^{(3)}$	3	3	-1	3	3	-1	1	1	1	1	-1	-1	-1	3	3	0	0	0	0	1	1	-1	-1
$\chi_\delta^{(3)}$	3	3	-1	3	3	-1	1	1	1	1	-1	3	3	-1	-1	0	0	0	0	-1	-1	1	1
$\chi_\alpha^{(2)}$	2	-2	0	2	-2	0	$-\sqrt{2}$	$\sqrt{2}$	$\sqrt{2}$	$-\sqrt{2}$	0	2	-2	0	0	1	-1	1	-1	0	0	$\sqrt{2}$	$-\sqrt{2}$
$\chi_\beta^{(2)}$	2	-2	0	2	-2	0	$\sqrt{2}$	$-\sqrt{2}$	$-\sqrt{2}$	$\sqrt{2}$	0	2	-2	0	0	1	-1	1	-1	0	0	$-\sqrt{2}$	$\sqrt{2}$
$\chi_\gamma^{(2)}$	2	-2	0	-2	2	0	$-\sqrt{2}$	$\sqrt{2}$	$-\sqrt{2}$	$\sqrt{2}$	0	0	0	-2	2	1	-1	-1	1	$\sqrt{2}$	$-\sqrt{2}$	0	0
$\chi_\delta^{(2)}$	2	-2	0	-2	2	0	$\sqrt{2}$	$-\sqrt{2}$	$\sqrt{2}$	$-\sqrt{2}$	0	0	0	-2	2	1	-1	-1	1	$-\sqrt{2}$	$\sqrt{2}$	0	0
$\chi_\alpha^{(4)}$	4	-4	0	4	-4	0	0	0	0	0	0	4	-4	0	0	-1	1	-1	1	0	0	0	0
$\chi_\beta^{(4)}$	4	-4	0	-4	4	0	0	0	0	0	0	0	0	-4	4	-1	1	1	-1	0	0	0	0
$\chi_\alpha^{(6)}$	6	-6	0	6	-6	0	$-\sqrt{2}$	$\sqrt{2}$	$\sqrt{2}$	$-\sqrt{2}$	0	-2	2	0	0	0	0	0	0	0	0	$-\sqrt{2}$	$\sqrt{2}$
$\chi_\beta^{(6)}$	6	-6	0	6	-6	0	$\sqrt{2}$	$-\sqrt{2}$	$-\sqrt{2}$	$\sqrt{2}$	0	-2	2	0	0	0	0	0	0	0	0	$\sqrt{2}$	$-\sqrt{2}$
$\chi_\gamma^{(6)}$	6	-6	0	-6	6	0	$-\sqrt{2}$	$\sqrt{2}$	$-\sqrt{2}$	$\sqrt{2}$	0	0	0	2	-2	0	0	0	0	$-\sqrt{2}$	$\sqrt{2}$	0	0
$\chi_\delta^{(6)}$	6	-6	0	-6	6	0	$\sqrt{2}$	$-\sqrt{2}$	$\sqrt{2}$	$-\sqrt{2}$	0	0	0	2	-2	0	0	0	0	$\sqrt{2}$	$-\sqrt{2}$	0	0